

TABLE 17. Expanded Estimates of Numbers, Percent, and Origin of Coded-wire Tagged (CWT) and Constant Fractional Marking Program (CFMP) Hatchery Fall-run Chinook Salmon Straying into the South Fork Trinity River (SFTR) Basin above Sandy Bar, 1984 through 1990 seasons (nd = no data).

CWT and CFMP Release Site:	year						
	1984	1985	1986	1987	1988	1989	1990
	Trinity River Basin						
Trinity River Hatchery (TRH)	(nd)	18 (10.2%)	9 (3.4%)	79 (17.4%)	18 (4.9%)	0 (0.0%)	0 (0.0%)
TRH Downstream Release	(nd)	2 (1.1%)	0 (0.0%)	4 (0.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Local Hatcheries a/	(nd)	2 (1.1%)	0 (0.0%)	11 (2.4%)	37 (10.1%)	5 (9.6%)	9 (4.0%)
RV Fin Clip (TRH)	3 (4.1%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Subtotal	3 (4.1%)	22 (12.4%)	9 (3.4%)	94 (20.7%)	55 (15.0%)	5 (9.6%)	9 (4.0%)
	Klamath River Basin						
IGH Downstream Release b/	(nd)	17 (9.7%)	1 (0.4%)	1 (0.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
LP Fin Clip (off-site)	0 (0.0%)	3 (1.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Subtotal	0 (0.0%)	20 (11.4%)	1 (0.4%)	1 (0.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
	Origin Unknown						
	18 (24.7%)						
<hr/>							
Numbers and Percent of Hatchery-origin							
Fish Captured at SFTR Weir	21 (28.8%)	42 (23.8%)	10 (3.8%)	95 (21.0%)	55 (15.0%)	5 (9.6%)	9 (4.0%)
Number of Fish trapped	73	176	264	455	368	52	223

a/ Includes Hoopa Valley Tribe hatcheries, Bureau of Indian Affairs rearing pond at Cappell Creek, and the Horse Linto Creek Hatchery.

b/ IGH = Iron Gate Hatchery